

ISSN 1682-8356
ansinet.org/ijps



INTERNATIONAL JOURNAL OF POULTRY SCIENCE

ANSI*net*

308 Lasani Town, Sargodha Road, Faisalabad - Pakistan
Mob: +92 300 3008585, Fax: +92 41 8815544
E-mail: editorijps@gmail.com

Effects of Age, Vitamin D3 and Fructooligosaccharide on Correlation of Different Bone Parameters in Broiler Chickens

A. Regassa¹ and W.K. Kim²

¹Department of Animal Science, University of Manitoba, Winnipeg, Manitoba, Canada

²Department of Poultry Science, University of Georgia, Athens, Georgia, USA

Abstract: Effects of age, vitamin D3 (VitD3) and Fructooligosaccharide (FOS) on the correlations of bone parameters were examined in broiler chickens. A total of 36 d-old male chicks were placed in battery brooders and fed a corn/soybean starter diet for 7 days and were randomly assigned to 12 pens each containing 3 birds. In group 1, birds were fed a corn/soybean starter diet and sacrificed on d 14 and 21. In groups 2 and 3, birds were fed a starter diet +250 µg VitD3/kg of diet and 4% FOS/kg of diet for 21 d, respectively. Birds were sacrificed to collect femur bones for measurement of bone parameters. Data were analyzed using Spearman's correlation analysis. Total mineral density (BMD) and trabecular density ($r_{14} = 0.99$, $p_{14} = 0.0001$; $r_{21} = 0.90$, $p_{21} = 0.0001$), total bone area and trabecular area ($r_{14} = 0.99$, $p_{14} = 0.0001$; $r_{21} = 1.00$, $p_{21} < 0.0001$) and trabecular mineral content and cortical mineral content ($r_{14} = 0.99$, $p_{14} < 0.0001$; $r_{21} = 0.96$, $p_{21} < 0.0001$) were strongly correlated both on d 14 and 21. In birds treated with VitD3 and FOS, bone mineral content (BMC) and cortical bone mineral content ($r = 0.95$, $p < 0.0001$) and BMD and trabecular mineral content ($r = 0.92$, $p = 0.0004$), were strongly correlated but most of the parameters were moderately or weakly correlated. In conclusion, results of this study show that age of birds, VitD3 and FOS treatments influence the relationships among the bone parameters.

Key words: Broiler, bone parameters, correlation

INTRODUCTION

In broilers, fast growth rates are generally correlated with musculoskeletal weakness because the development of bone in these animals is not well organized, less dense and more porous than slow growing ones (Williams *et al.*, 2004; Bennett, 2008). This is attributed to the imbalance between meat production and skeletal growth caused by genetic selection resulting in abnormal skeletal development (Oviedo-Rondon *et al.*, 2006). Such skeletal abnormalities are among the problems facing the poultry industry. For instance, leg problems and skeletal disorders, such as lameness, twisted legs, tibial dyschondroplasia and crooked toes, are the common causes of culling and mortality in broiler chicks (Bradshaw *et al.*, 2002; Oviedo-Rondon *et al.*, 2006).

Vitamin D3 (VitD3) can be synthesized under skin through ultraviolet irradiation of 7-dehydrocholesterol or absorbed from the diet in the intestinal tract (DeLuca, 2004). It is biologically inactive in this form and needs to be converted to 25-hydroxy vitamin D3 in the liver and subsequently to 1,25-dihydroxy vitamin D3 in the kidney (DeLuca, 2004). The hormone form of VitD3, 1,25-dihydroxy vitamin D3 acts through a nuclear receptor to express its numerous functions, including bone homeostasis, calcium and phosphate absorption in the small intestine, calcium mobilization in bone and

calcium reabsorption in the kidney (Leeson and Summers, 2001; DeLuca *et al.*, 2004). High levels of VitD3 reduces incidence of leg problem, including tibial dyschondroplasia and rickets in broilers (Whitehead *et al.*, 2004; Atencio *et al.*, 2005; Driver *et al.*, 2006) and increasing dietary vitamin D3 improves the walking ability and welfare status of broiler chickens reared at high stocking densities (Sun *et al.*, 2013). Additionally, high levels of VitD3 have been suggested to increase bone growth and mineral deposition in broiler chicks (Kim *et al.*, 2011).

Fructooligosaccharides (FOS) are low-molecular-weight non-digestible carbohydrates that have the potential to increase bone formation and reduce bone resorption (Ohta *et al.*, 1995; Takahara *et al.*, 2000). Studies in rats have demonstrated that FOS enhances true Ca, Mg, Fe and P absorptions in the intestine (Ohta *et al.*, 1995; Kashimura *et al.*, 1996; Morohashi *et al.*, 1998), can increase femoral bone volume and mineral concentrations (Takasaki *et al.*, 2000), femoral calcium content and bone mineral density (Zafar *et al.*, 2004) and bone mass and biomechanical properties (Lobo *et al.*, 2006). Studies have shown that FOS have the potential to maintain bone strength during molting of white leghorn hens (Kim *et al.*, 2006) but do not have any beneficial effects on bone growth and skeletal integrity in growing male broiler chicks (Kim *et al.*, 2011).

Studies have been conducted to examine bone mineralization rates and growth in different chicken types. For instance, differences in mineralization rates and growth characteristics among different skeletal structures have been reported in commercial broiler lines (Applegate and Lilburn, 2002). Lower epiphyseal ash values at younger ages and smaller body weights in 2010 commercial strains of Pekin ducks suggest the need to critically monitor the factors that influence bone mineralization (van Wyhe *et al.*, 2012). The results of a study in commercial male and female broilers have indicated that life cycle changes do not significantly affect bone mineralization and bone size traits (Talaty *et al.*, 2009). Large differences in bone traits have been reported in several purebred lines of meat-type chickens suggesting the potential to genetically select birds for increased bone mineral density (BMD) (Korver *et al.*, 2004). The study also showed changes in BMD of individual birds can be measured over time and BMD at specific time points can be correlated with production parameters and eggshell quality traits (Korver *et al.*, 2004).

Studies have also been conducted to establish the importance of some bone parameters in predicting the health of bone. Assessment of cortical bone strength has been suggested to predict the risk of bone fracture or as a choice of suitable therapeutic strategies in orthopedic surgery (Augat *et al.*, 1996, 1998). Similarly, BMD has been reported to be helpful for the estimation of bone strength and prediction of intracortical porosity and parameters related to porous structures of cortical bone (Wachter *et al.*, 2002).

The degrees of associations among bone parameters and production traits have also been examined in chicken. High correlation coefficients have been found for BMD and bone mineral content (BMC) between live and excised tibial bones in White Leghorn hens (Schreiweis *et al.*, 2005). Correlation between third-toe-amplitude dependent speed of sound measurement, shear strength and radiographic density values have been reported in caged and free-range hens (Fleming *et al.*, 2004). However, absence of correlation between egg production or egg quality traits and bone quality measures have been reported in White Leghorns housed in conventional, modified and commercially available colony battery cages (Jendral *et al.*, 2008).

These works have focused on the estimation of the degree of correlation between bone parameters and production traits or the effects of dietary treatments on the magnitude of bone parameter itself but not on the degree of correlation among different bone parameters. Thus the degree of association among different bone parameters due to age difference and dietary treatments, such as vitD3 and FOS, on the correlation of these parameters are largely unknown. Therefore, this study was conducted to determine the effects of age

(Days 14 and 21), dietary treatments (VitD3 and FOS) on the degree of correlation among several bone parameters in broilers.

MATERIALS AND METHODS

General procedures: A total of 36 day-old male broiler chicks were placed in heated Petersime battery brooders and fed a corn/soybean broiler starter diet for seven days. Then, the chicks were individually weighed and randomly assigned to 12 pens of three birds each. The starter diet contained 23% crude protein, 3,200 kcal/kg metabolic energy, 1% calcium, 0.45% available phosphorus and 90 µg/kg VitD3. In treatment one, where we examined how bone parameters correlate in a normal corn/soybean dietary condition, birds were sacrificed on Days 14 and 21. In treatments two and three, birds were fed a control corn/soybean meal diet with 250 µg VitD3/kg of diet and 4% FOS for 21 days, respectively. At the end of each time points, birds were sacrificed by cervical dislocation to collect femur bones for peripheral quantitative computed tomography (pQCT) and mechanical testing. In order to determine how age affects the correlation among bone parameters in chicks fed a normal corn/soybean meal diet, the data of different bone parameters on Days 14 and 21 were analyzed separately for their respective correlation coefficients. To examine the effects of VitD3 and FOS on the correlation of bone parameters, data of each group were analyzed separately for their respective correlation coefficients. All animal procedures were done according to the University committee guideline for Animal Care.

Peripheral quantitative computed tomography: Bone samples were scanned using a XCT Research instrument (Stratec, Norland, Fort Atkinson, WI). This model has a minimum voxel size of 0.07 mm and a scanning beam thickness of 0.50 mm. Scan sites included the mid-shaft femurs (3 slices of each bone located at one-half the total bone length ± 2 mm) for volumetric bone mineral density (vBMD) of the cortical bones the proximal femurs were scanned to assess metaphyseal trabecular vBMD (3 slices located 8.0, 8.5 and 9.0 mm from the proximal end of the bone). All scans were obtained at a scan speed of 2.5 mm/sec, with a voxel resolution of 0.07x0.07x0.50 mm. In addition, the mid diaphyseal cross-sectional moment of inertia (CSMI), a measure of bone material in a cross section area and indicator of its distribution around the neutral bending axis to resist bending (Bennett, 2008), was obtained with respect to the neutral bending axis of the femur bone shaft during 3-point bending (mechanical property testing).

Mechanical testing: After pQCT scans were completed, mechanical properties of the mid shaft femurs were determined by 3-point bending to failure with an Instron

1125 servo-controlled testing machine (Instron Corp., Canton, MA) according to previously published procedures (Allen and Bloomfield, 2003). The femurs were thawed at room temperature and placed posterior side down on metal pin supports located ± 10 mm (femurs) from the mid diaphysis testing site. With a 1,000 Lb (455 kg) load cell, quasi-static loading (2.5 mm/min) was applied to the anterior surface of the femurs until fractures occurred. All specimens were sprayed with PBS just before testing to maintain hydration. Displacements were monitored by a linear variable differential transformer interfaced with a personal computer. Raw data, collected at 10 Hz as load versus displacement curves, were analyzed with Table-Curve 2.0 software (Jandel, San Rafael, CA). Structural variables (ultimate load and stiffness) were obtained directly from the load: displacement curves. The maximum load obtained was defined as the ultimate load (UL, in N) and the slope of the elastic portion of the curve was defined as stiffness (S, in N/mm). Bone tissue material properties were calculated by normalizing structural properties for cross-sectional bone geometry at the site of testing by using CSMI (in mm^4 from pQCT), bone diameter (D, in mm) as measured by calipers and the appropriate bottom support span distance (L, which was 20 or 60 mm). The appropriate formulas for elastic modulus (E, in GPa) and ultimate stress (US, in MPa) were as follows:

$$E = S \times L^3 / (48,000 \times \text{CSMI}) \text{ and } US = UL \times L \times D / (8 \times \text{CSMI})$$

Statistical analysis: Bone parameter data were analyzed using Spearman's correlation analysis (SAS Institute, 2008). Correlations among the parameters were considered significant at $p < 0.05$. Strongly correlated parameters were identified using $r = 0.90$.

RESULTS AND DISCUSSION

Correlation analysis of bone parameters in chicks on Days 14 and 21: Age based correlation analysis of bone parameters in birds fed a normal corn/soybean meal diet revealed that most of the parameters were strongly correlated with each other both on Day 14 and 21 with slightly declined or weaker correlations on Day 21 (Fig. 1 and Table 1 and 2). Parameters that were strongly correlated both on Day 14 and 21 include: total mineral density with trabecular density ($r_{14} = 0.98$, $p_{14} < 0.0001$; $r_{21} = 0.91$, $p_{21} = 0.0001$), total bone area with trabecular area ($r_{14} = 0.99$, $p_{14} = 0.0001$; $r_{21} = 1.00$, $p_{21} < 0.0001$), trabecular mineral content with cortical mineral content ($r_{14} = 0.99$, $p_{14} < 0.0001$; $r_{21} = 0.96$, $p_{21} < 0.0001$), total mineral content with trabecular mineral content ($r_{14} = 0.99$, $p_{14} < 0.0001$; $r_{21} = 0.96$, $p_{21} < 0.0001$), total bone area with cortical area ($r_{14} = 0.97$, $p_{14} = 0.0001$; $r_{21} = 0.97$, $p_{21} < 0.0001$) and cortical mineral content with cortical area ($r_{14} = 0.97$, $p_{14} = 0.0001$; $r_{21} = 0.97$, $p_{21} < 0.0001$).

These results suggest that there is a trend where total bone parameters, such as total bone area, total BMD, total BMC, are strongly correlated with their respective regional parameters, such as trabecular and cortical area, trabecular mineral density, trabecular and cortical mineral content (Table 1 and 2). This can be attributed to the fact that regional bone parameters are the components of the total parameters (Franck and Munz, 2000). Bone mineral density measurements have been suggested to be combined with trabecular micro-architecture to predict the fracture risk and the severity of osteoporosis in middle-aged men showing the association between the two parameters (Legrand *et al.*, 2000). Additionally, significant correlation between cortical and trabecular bone mineral content has previously been reported in women (Bohr and Schaadt, 1985).

Mechanical strength is highly correlated with stiffness, particularly on Day 14 ($r = 0.90$, $p = 0.0009$) (Fig. 2). Mechanical strength of a material is its ability to withstand an applied stress without forming a fracture (Duane, 2003; WuBian *et al.*, 2010) and stiffness is the resistance of an elastic body to deformation (Gopalakrishnan and Zukoski, 2007). Their strong correlation in this study can be explained by the fact that both of them are the factors influencing bone strength and measuring one of these parameters could provide an indirect value for the other.

The correlations between mechanical strength and stiffness ($r_{14} = 0.90$, $p_{14} < 0.0009$; $r_{21} = 0.74$, $p_{21} = 0.022$) (Fig. 2), total bone area with cortical thickness ($r_{14} = 0.88$, $p_{14} = 0.0038$; $r_{21} = 0.80$, $p_{21} = 0.0095$), trabecular mineral content ($r_{14} = 0.87$, $p_{14} = 0.0051$; $r_{21} = 0.79$, $p_{21} = 0.01$), trabecular mineral content with trabecular area ($r_{14} = 0.87$, $p_{14} = 0.0051$; $r_{21} = 0.79$, $p_{21} = 0.01$), trabecular density with cortical thickness ($r_{14} = 0.89$, $p_{14} = 0.0024$; $r_{21} = 0.71$, $p_{21} = 0.03$) and trabecular mineral content ($r_{14} = 0.94$, $p_{14} = 0.0005$; $r_{21} = 0.75$, $p_{21} = 0.02$) showed declining tendencies with age (Table 1 and 2). Previously, an age-related decline in trabecular density has been reported in a human study (Singer and Breidahl, 1990). This age related declining correlations between the parameters in the present study might indicate that the parameters are not changing proportionally as age increases and this requires further investigation.

Although most of the parameters showed strong to moderate correlation at both stages, some of them were poorly correlated on day 21 (Fig. 3 and Table 2). Parameters that were poorly correlated on Day 21 include BMD with cortical area ($r_{21} = 0.1$, $p_{21} = 0.79$), cortical thickness ($r_{21} = 0.46$, $p_{21} = 0.2$), cortical mineral content ($r_{21} = 0.3$, $p_{21} = 0.42$) and trabecular mineral content ($r_{21} = 0.49$, $p_{21} = 0.17$) (Table 2). These data are consistent with those from human studies.

Table 1: Correlation coefficients and their respective probabilities of bone parameters obtained from birds fed starter diet for 14 days

| | Trabecular area (mm ²) | Cortical area (mm ²) | Cortical thickness (mm) | Total mineral content (g/cm) | Trabecular mineral content (g/cm) | Cortical mineral content (g/cm ³) |
|---|------------------------------------|----------------------------------|-------------------------|------------------------------|-----------------------------------|---|
| Total area (mm ²) | r = 0.99, p = 0.0001 | r = 0.97, p = 0.0001 | r = 0.88, p = 0.0038 | r = 0.91, p = 0.0016 | r = 0.87, p = 0.0051 | r = 0.91, p = 0.0016 |
| Cortical area (mm ²) | r = 0.97, p = 0.0001 | 1.0000 | r = 0.97, p = 0.0001 | r = 0.97, p = 0.0001 | r = 0.95, p = 0.0003 | r = 0.97, p = 0.0001 |
| Trabecular area (mm ²) | 1.0000 | r = 0.97, p = 0.0001 | r = 0.88, p = 0.0039 | r = 0.91, p = 0.0016 | r = 0.87, p = 0.0051 | r = 0.91, p = 0.0016 |
| Trabecular mineral content (g/cm) | r = 0.87, p = 0.0051 | r = 0.95, p = 0.0003 | r = 0.97, p = 0.0001 | r = 0.99, p = 0.0001 | 1.0000 | r = 0.99, p = 0.0001 |
| Total mineral density (mg/cm ³) | r = 0.1, p = 0.79 | r = 0.76, p = 0.028 | r = 0.86, p = 0.0064 | r = 0.88, p = 0.0038 | r = 0.92, p = 0.0013 | r = 0.87, p = 0.004 |
| Trabecular density (mg/cm ³) | r = 0.2, p = 0.62 | r = 0.80, p = 0.016 | r = 0.89, p = 0.0024 | r = 0.89, p = 0.0028 | r = 0.94, p = 0.0005 | r = 0.89, p = 0.003 |

r = correlation coefficient, p = probability

Table 2: Correlation coefficients and their respective probabilities of bone parameters obtained from birds fed starter diet for 21 days

| | Trabecular area (mm ²) | Cortical area (mm ²) | Cortical thickness (mm) | Total mineral content (g/cm) | Trabecular mineral content (g/cm) | Cortical mineral content (g/cm) |
|---|------------------------------------|----------------------------------|-------------------------|------------------------------|-----------------------------------|---------------------------------|
| Total area (mm ²) | r = 1.000, p < 0.0001 | r = 0.97, p = 0.0001 | r = 0.80, p = 0.0095 | r = 0.91, p = 0.0005 | r = 0.79, p = 0.01 | r = 0.91, p = 0.0006 |
| Cortical area (mm ²) | r = 0.96, p < 0.0001 | 1.0000 | r = 0.92, p = 0.0003 | r = 0.98, p = 0.0001 | r = 0.91, p = 0.0005 | r = 0.97, p = 0.0001 |
| Trabecular area (mm ²) | 1.0000 | r = 0.96, p < 0.0001 | r = 0.80, p = 0.0095 | r = 0.91, p = 0.0005 | r = 0.79, p = 0.01 | r = 0.91, p = 0.0006 |
| Trabecular mineral content (g/cm) | r = 0.79, p = 0.01 | r = 0.91, p = 0.0005 | r = 0.98, p = 0.0005 | r = 0.96, p = 0.0001 | 1.0000 | r = 0.96, p = 0.0001 |
| Total mineral density (mg/cm ³) | r = 0.1, p = 0.79 | r = 0.14, p = 0.70 | r = 0.46, p = 0.2 | r = 0.3, p = 0.42 | r = 0.49, p = 0.17 | r = 0.30, p = 0.42 |
| Trabecular density (mg/cm ³) | r = 0.19, p = 0.62 | r = 0.43, p = 0.24 | r = 0.71, p = 0.03 | r = 0.55, p = 0.12 | r = 0.75, p = 0.02 | r = 0.54, p = 0.13 |

r = correlation coefficient, p = probability

Table 3: Correlation coefficients and their respective probabilities of bone parameters obtained from birds treated with 250 µg of vitamin D3 per kilogram of diet for 21 days

| | Ultimate strength (N/cm ²) | Cortical area (mm ²) | BMC (mm) | Trabecular mineral density (mg/cm ³) | Cortical mineral content (mm) |
|---------------------------------------|--|----------------------------------|----------------------|--|-------------------------------|
| Mechanical force (N/cm ²) | r = 0.83, p = 0.005 | r = 0.52, p = 0.14 | r = 0.65, p = 0.05 | r = 0.74, p = 0.02 | r = 0.68, p = 0.04 |
| BMD (mg/cm ³) | r = 0.78, p = 0.01 | r = 0.59, p = 0.08 | r = 0.67, p = 0.04 | r = 0.95, p < 0.0001 | r = 0.70, p = 0.03 |
| Total bone area (mm ²) | r = -0.88, p = 0.0016 | r = 0.85, p = 0.0032 | r = 0.83, p = 0.005 | r = 0.13, p = 0.73 | r = 0.80, p = 0.0087 |
| Cortical area (mm ²) | r = 0.06, p = 0.87 | - | r = 0.97, p < 0.0001 | r = 0.60, p = 0.08 | r = 0.97, p < 0.0001 |
| Trabecular area (mm ²) | r = -0.30, p = 0.41 | r = 0.85, p = 0.003 | r = 0.83, p = 0.005 | r = 0.13, p = 0.73 | r = 0.8, p = 0.007 |

r = correlation coefficient, p = probability

Table 4: Correlation coefficients and their respective probabilities of bone parameters obtained from birds treated with 4% of fructo oligosaccharide per kilogram of diet for 21 days

| | Ultimate strength (N/mm ²) | Mechanical force (N/mm ²) | Total area (mm ²) | Cortical thickness (mm) | Trabecular mineral content (g/cm ³) | Cortical mineral content (g/cm) |
|------------------------------------|--|---------------------------------------|-------------------------------|-------------------------|---|---------------------------------|
| Trabecular area (mm ²) | r = -0.88, p = 0.0016 | r = -0.87, p = 0.0022 | r = 1.000, p < 0.0001 | r = 0.91, p = 0.0005 | r = 0.7, p = 0.03 | r = 0.84, p = 0.004 |
| Cortical area (mm ²) | r = 0.72, p = 0.02 | r = -0.83, p = 0.0058 | r = 0.91, p = 0.0005 | r = 0.91, p = 0.0005 | r = 0.89, p = 0.0012 | r = 0.86, p = 0.0025 |
| BMC (g/cm ³) | r = -0.54, p = 0.13 | r = 0.39, p = 0.29 | r = 0.86, p = 0.003 | r = 0.86, p = 0.0031 | r = 0.92, p = 0.0004 | r = 0.99, p < 0.0001 |
| Cortical thickness (mm) | r = -0.38, p = 0.30 | r = 0.36, p = 0.33 | r = 0.63, p = 0.06 | r = 1.000 | r = 0.87, p = 0.0021 | r = 0.80, p = 0.0091 |

r = correlation coefficient, p = probability




Fig. 1(a-b): Correlation between total mineral density and trabecular density on day 14 (a) and day 21 (b) of the experimental period. Although showing a declining tendency with age, the parameters were strongly correlated both on day 14 ($r = 0.98092$, $p < 0.0001$) and day 21 ($r = 0.90964$, $p = 0.0001$)




Fig. 2(a-b): Correlation between mechanical strength and bone stiffness on day 14 (a) and day 21 (b) of the experimental period. The parameters were strongly correlated on day 14 ($r = 0.90$, $p = 0.0009$) but moderately correlated on day 21 ($r = 0.74$, $p = 0.0225$)

Age-dependent decline in cortical BMC and trabecular BMD at the distal radius, spinal BMD, trabecular bone volume fraction, trabecular thickness, trabecular number and fractal dimension have been reported in premenopausal women (Majumdar *et al.*, 1997). An increased growth rate is associated with lower mineral density in broiler chicken (Leterrier and Nys, 1992) and consistent with some of these data, the correlations of some of the parameters were weaker on day 21 than on day 14. On the other hand Lee *et al.* (2001) have reported a strong negative correlation between total and regional BMD in menopausal Japanese women showing the effect of interspecies differences on the correlation of bone parameters.

Correlation analysis of bone parameters in Vit D3 and FOS fed birds: As opposed to a normal corn/soybean meal dietary condition where most parameters were strongly correlated, VitD3 and FOS treatments affected the correlation of the parameters. Except for strong

correlation between BMC and cortical bone mineral content ($r = 0.95$, $p < 0.0001$), BMD and trabecular mineral content ($r = 0.92$, $p = 0.0004$), mechanical strength and BMD ($r = 0.89$, $p = 0.0012$), trabecular area and cortical thickness ($r = 0.91$, $p = 0.0005$) and cortical area and total area ($r = 0.91$, $p = 0.0005$), most of the parameters were moderately or weakly correlated in birds treated with VitD3 and FOS (Table 3 and 4, Fig. 4). There were also parameters which were negatively correlated emphasizing the effects of these dietary treatments on the association of bone parameters (Table 3 and 4). Consistent with strong correlation between BMD and mechanical strength in the present study, high correlation coefficient between BMD and strength has been reported in female cadavers indicating the dependence of bone strength on BMD (Graham *et al.*, 2007). On the other hand, low correlation coefficients have been reported between shear force and BMD or BMC in three weeks old broiler chicks received adequate, medium and low dietary Ca and P levels




Fig. 3(a-d): Correlation between BMD and BMC and between BMD and trabecular mineral content on day 14 (a and c) and day 21 (b and d) of the experimental period. The parameters were strongly correlated on day 14 ($r_a = 0.88$, $p_a = 0.0038$, $r_c = 0.91$, $p_c = 0.0013$) but did not correlate on day 21 ($r_b = 0.30$, $p_b = 0.42$, $r_d = 0.49$, $p_d = 0.1758$)




Fig. 4(a-b): Correlation between mechanical force and BMD in birds treated with 4% FOS (a) and between mechanical force and ultimate strength in birds treated with 250 µg of VitD3/Kg of diet (b) r_a , r_b , r_c and r_d stand for the coefficients of correlation between the parameters depicted on figure 3A, B, C and D, respectively and p_a , p_b , p_c and p_d stand for their respective probabilities

(Onyango *et al.*, 2003) suggesting a possible interspecies variation among different bone parameters.

Conclusion: In conclusion, the results of this study showed that most of the parameters in broiler chicks fed a normal corn/soybean meal diet are strongly and

moderately correlated, and age of the birds appears to influence the correlation of bone parameters providing an insight into how the parameters are associated during the broiler growth period and dietary supplementation of VitD3 and FOS affects the direction and magnitude of correlation.

REFERENCES

- Allen, M.R. and S.A. Bloomfield, 2003. Hind limb unloading has a greater effect on cortical compared with cancellous bone in mature female rats. *J. Appl. Physiol.*, 94: 642-650.
- Applegate, T.J. and M.S. Lilburn, 2002. Growth of the femur and tibia of a commercial broiler line. *Poult. Sci.*, 81: 1289-1294.
- Atencio, A., H.M. Edwards and G.M. Pesti, 2005. Effect of the level of cholecalciferol supplementation of broiler breeder hen diets on the performance and bone abnormalities of the progeny fed diets containing various levels of calcium or 25-hydroxycholecalciferol. *Poult. Sci.*, 84: 1593-1603.
- Augat, P., B. Fan and N.E. Lane, 1998. Assessment of bone mineral at appendicular sites in females with fractures of the proximal femur. *Bone*, 22: 395-402.
- Augat, P., H. Reeb and L.E. Claes, 1996. Prediction of fracture load at different skeletal sites by geometric properties of the cortical shell. *J. Bone. Miner. Res.*, 11: 1356-1363.
- Bennett, M.B., 2008. Post-hatch growth and development of the pectoral and pelvic limbs in the black noddly, *Anous minutus*. *Comp. Biochem. Physiol. Part A.*, 150: 159-168.
- Bohr, H. and O. Schaadt, 1985. Bone mineral content of the femoral neck and shaft: relation between cortical and trabecular bone. *Calcif. Tissue. Int.*, 37: 340-344.
- Bradshaw, R.H., R.D. Kirkden and D.M. Broom, 2002. A review of the aetiology and pathology of leg weakness in broilers in relation to welfare. *Avian. Poul. Biol. Rev.*, 13: 45-103.
- DeLuca, H.F., 2004. Overview of general physiologic features and functions of vitamin D. *Am. J. Clin. Nutr.*, 80: 1689-1696.
- Driver, J.P., A. Atencio, G.M. Pesti, H.M. Edwards and R.I. Bakalli, 2006. The effect of maternal dietary vitamin D3 supplementation on performance and tibial dyschondroplasia of broiler chicks. *Poult. Sci.*, 85: 39-47.
- Duane, K., 2003. *Fundamentals of Biomechanics*. Kluwer Academic/Plenum Publishers, New York.
- Fleming, R.H., D. Korver, H.A. McCormack and C.C. Whitehead, 2004. Assessing bone mineral density in vivo: digitized fluoroscopy and ultrasound. *Poult. Sci.*, 83: 207-214.
- Franck, H. and M. Munz, 2000. Total body and regional bone mineral densitometry and soft tissue measurements: Correlations of bone mineral density parameter to lumbar spine and hip. *Calcif. Tissue. Int.*, 67: 111-115.
- Gopalakrishnan, V. and C.F. Zukoski, 2007. Delayed flow in thermo-reversible colloidal gels. *J. Rheol.*, 51: 623-644.
- Graham, J., A. Chul, H. Nabila and D.B. Barbara, 2007. Effect of bone density on vertebral strength and stiffness after percutaneous vertebroplasty. *SPINE*, 32: 505-511.
- Jendral, M.J., D.R. Korver, J.S. Church and J.J. Feddes, 2008. Bone mineral density and breaking strength of white leghorns housed in conventional, modified and commercially available colony battery cages. *Poult. Sci.*, 87: 828-837.
- Kashimura, J., M. Kimura and Y. Itokawa, 1996. The effects of isomaltulose, isomalt and isomaltulose-based oligomers on mineral absorption and retention. *Biol. Trace. Elem. Res.*, 54: 239-250.
- Kim, W.K., S.A. Bloomfield and S.C. Ricke, 2011. Effects of age, vitamin D3 and fructooligosaccharides on bone growth and skeletal integrity of broiler chicks. *Poult. Sci.*, 90: 2425-2432.
- Kim, W.K., L.M. Donalson, A.D. Mitchell, L.E. Kubena, D.J. Nisbet and S.C. Ricke, 2006. Effects of alfalfa and fructooligosaccharide on molting parameters and bone qualities using dual energy X-ray absorptiometry and conventional bone assays. *Poult. Sci.*, 85: 15-20.
- Korver, D.R., J.L. Saunders-Blades and K.L. Nadeau, 2004. Assessing bone mineral density *in vivo*: quantitative computed tomography. *Poult. Sci.*, 83: 222-229.
- Lee, J.S., K. Kawakubo, H. Sato, Y. Kobayashi and Y. Haruna, 2001. Relationship between total and regional bone mineral density and menopausal state, body composition and life style factors in overweight Japanese women. *Int. J. Obes.*, 25: 880-886.
- Leeson, S. and J.D. Summers, 2001. *Scott's Nutrition of the Chicken*. Publ Univ Books, Guelph, Ontario, Canada.
- Légrand, E., D. Chappard, C. Pascaretti, M. Duquenne, S. Krebs, V. Rohmer, M.F. Basle and M. Audran, 2000. Trabecular bone microarchitecture, bone mineral density and vertebral fractures in male osteoporosis. *J. Bone. Miner. Res.*, 15: 13-19.
- Leterrier, C. and Y. Nys, 1992. Composition, cortical structure and mechanical properties of chicken tibiotarsi: Effect of growth rate. *Br. Poult. Sci.*, 33: 925-939.
- Lobo, A.R., C. Colli and T.M. Filisetti, 2006. Fructooligosaccharides improve bone mass and biomechanical properties in rats. *Nutr. Resea.*, 26: 413-420.
- Majumdar, S., H.K. Genant, S. Grampp, D.C. Newitt, V.H. Truong, J.C. Lin and A. Mathur, 1997. Correlation of trabecular bone structure with age, bone mineral density and osteoporotic status: In vivo studies in the distal radius using high resolution magnetic resonance imaging. *J. Bone. Miner. Res.*, 12: 111-118.

- Morohashi, T., T. Sano, A. Ohta and S. Yamada, 1998. True calcium absorption in the intestine is enhanced by fructooligosaccharide feeding in rats. *J. Nutr.*, 128: 1815-1818.
- Ohta, A.M., O.S. Baba, T. Adachi T, Sakata and E. Sakaguchi, 1995. Calcium and magnesium absorption from the colon and rectum are increased in rats fed fructooligosaccharides. *J. Nutr.*, 125: 2417-2424.
- Onyango, E.M. P.Y. Hester, R. Strohshine and O. Adeola, 2003. Bone densitometry as an indicator of percentage tibia ash in broiler chicks fed varying dietary calcium and phosphorus levels. *Poult. Sci.*, 82: 1787-1791.
- Oviedo-Rondon, E.O., P.R. Ferket and G.B. Havenstein, 2006. Understanding long bone development in broilers and turkeys. *Avian. Poult. Biol. Rev.*, 17: 77-88.
- SAS Institute, 2008. SAS System for Statistical Analysis. Release 9.2., SAS Inst., Inc., Cary, NC.
- Schreiweis, M.A., J.I. Orban, M.C. Ledur, D.E. Moody and P.Y. Hester, 2005. Validation of dual-energy X-ray absorptiometry in live white leghorns. *Poult. Sci.*, 84: 91-99.
- Singer, K.P. and P.D. Breidiah, 1990. Vertebral body trabecular density at the thoracolumbar junction using quantitative computed tomography. A post-mortem study. *Acta. Radiol.*, 31: 37-40.
- Sun, Z.W., L. Yan, G. YY, J.P. Zhao, H. Lin and Y.M. Guo, 2013. Increasing dietary vitamin D3 improves the walking ability and welfare status of broiler chickens reared at high stocking densities. *Poult. Sci.*, 92: 3071-3079.
- Takahara, S., T. Morohashi, T. Sano, A. Ohta, S. Yamada and R. Sasa, 2000. Fructooligosaccharide consumption enhances femoral bone volume and mineral concentrations in rats. *J. Nutr.*, 130: 1792-1795.
- Takasaki, M., H. Inaba, A. Ohta, Y. Motohashi, K. Sakai, H. Morris and K. Sakuma, 2000. Dietary short-chain fructooligosaccharides increase calbindin-D9k levels only in the large intestine in rats independent of dietary calcium deficiency or serum 1,25 dihydroxy vitamin D levels. *Int. J. Vitam. Nutr. Res.*, 70: 206-213.
- Talaty, P.N., M.N. Katanbaf and P.Y. Hester, 2009. Life cycle changes in bone mineralization and bone size traits of commercial broilers. *Poult. Sci.*, 88: 1070-1077.
- Van Wyhe, R.C., T.J. Applegate, M.S. Lilburn and D.M. Karcher, 2012. A comparison of long bone development in historical and contemporary ducks. *Poult. Sci.*, 91: 2858-2865.
- Wachter, N.J., G.D. Krischak, M. Mentze, M.R. Sarkar, T. Ebinger, L. Kinzl, L. Claes and L.P. Augat, 2002. Correlation of bone mineral density with strength and micro structural parameters of cortical bone *in vitro*. *Bone*, 31: 90-95.
- Whitehead, C.C., H.A. McCormack, L. McTeir and R.H. Fleming, 2004. High vitamin D3 requirements in broilers for bone quality and prevention of tibial dyschondroplasia and interactions with dietary calcium, available phosphorus and vitamin A. *Br. Poult. Sci.*, 45: 425-436.
- Williams, B.D., D.H. Waddington, C. Murray and C. Farquharson, 2004. Bone strength during growth: Influence of growth rate on cortical porosity and mineralization. *Calcif. Tissue. Int.*, 74: 236-245.
- WuBian, T., S. Zheng Ming, H. Hitoshi and D. YuLei, 2010. Micro structural evolution and mechanical properties of Ti3SiC2-TiC composites. *J. Alloys. Comp.*, 502: 49-53.
- Zafar, T., A.C. Weaver, Y. Zhao, B.R. Martin and M.E. Wastney, 2004. Non-digestible oligosaccharides increase calcium absorption and suppress bone resorption in ovariectomized rats. *J. Nutr.*, 134: 399-402.